

TECHNICAL DATA SHEET

LINDEN EXTRACT / A / HGL / CA11 /



Trade Name: LINDEN EXTRACT / A / HGL / CA11 /.

Product Code: ETIAHGLCA11.

INCI Name: AQUA, GLYCERIN, TILIA CORDATA EXTRACT, POTASSIUM SORBATE.

INCI Status: Conforms.

REACH Status: Low Volume Exemption.

CAS Number: 7732-18-5; 56-81-5; 84929-52-2; 24634-61-5 / 590-00-1.

EC Number: 231-791-2; 200-289-5; 284-536-2; 246-376-1.

Origin: GLYCERIN: Vegetable origin.

TILIA CORDATA EXTRACT: Vegetable origin. POTASSIUM SORBATE: Synthetic origin.

Processing: GMO Free. / No Ethoxylation. / No Irradiation. / No Sulphonation. / No BSE/TSE. / No CMR. / No SVHC. / No Nanomaterial. / No DEG. / No DEHP. / No Formaldehyde. / No Phthalates. /.

Additives: Preservatives: Potassium sorbate. Paraben Free. / Antioxidants: None. / Other additives: None.

Solvents Used: Water and glycerin.

Appearance: Clear or slightly opalescent liquid. **Colour:** Characteristic (yellowish to deep brown).

Odour: Characteristic.

Specific gravity: 1,050 - 1,300 g/cm³ (20°C approx.).

Direct pH value: 4,5 - 6,5 (20°C approx.).

Soluble/ Miscible: Water. Surfactants. Hydroalcoholic, hydroglyceric and hydroglycolic mixtures.

Ecological Information: Sustainable.

Microbial Count: Total aerobics: Max. 300 microorganisms / gram.
Fungi and yeasts: Max. 200 microorganisms / gram.

Pathogens: Total absence in 1 gram.

Suggested Use Levels: 2.0 - 10.0%.

Suggested Applications: Preparations to treat irritated, damaged, reddened and sensitive skin.

Properties: Emollient, calming, protective, softening, relaxing and anti-inflammatory.

Industrial packing: 1Kg., 5Kg., 25Kg.

Storage conditions: It is very important to keep the product in well closed containers, away from direct sunlight at temperatures not over 15°C. In case of not having refrigerators, store the product in the coolest and free from artificial light area.

Bibliography: - Atlas Ilustrado de Plantas Medicinales y Curativas, Ed. Susaeta.

- Dumont's grosse Kräuter Enzyklopädie.
- Fitocosmeticos ALVAREZ CRUZ i BAGUE SERRANO.
- Gran enciclopedia de las plantas medicinales, BERDONCES i SERRA.
- Plantas medicinales, FONT i QUER.
- Proserpio G., Martelli A., Patri GF.; Elementi di Fitocosmeci; Ed. Sepem / Milano 1983.
- Websites.

OVERVIEW

info@biogrundl.es

TECHNICAL DATA SHEET

Linden (Tilia cordata) is a tree which exists in many European countries, belonging to the tiliaceae family. In the Mediterranean countries it flowers between May and June, its flowers have a distinctive, penetrating aroma with calming and tranquilizer properties. From these flowers linden extract is obtained.

Its major active ingredient is essential oil, made up of hydrocarbons (C18-C30), phenylethanol, phenylethyl acetate, geraniol, geranyl acetate, eugenol, linalol and farnesol; in addition to flavonoids; coumarins; tannins; triterpenes; polyphenolic acids (caffeic, chlorogenic); saponins and mucilages.

USE AND DOSAGE

The incorporation of linden extract in cosmetics is justified by its emollient, calming, protective, softening, relaxing and anti-inflammatory properties.

Its use is very common in preparations to treat irritated, damaged, reddened and sensitive skin.

If turbidity is observed, it is advisable to previously dissolve the product in a portion of the water provided to manufacture the final product and filter after 12-24 hours.

The recommended concentrations of use are between 2% and 10%. For very special effects the dosage can raise up to 15%.

PHYSICAL AND CHEMICAL PROPERTIES

Appearance: Clear or slightly opalescent liquid. **Colour:** Characteristic (yellowish to deep brown).

Odour: Characteristic.

Direct pH value: 4,5 - 6,5 (20°C approx.).

Specific gravity: 1,050 - 1,300 g/cm³ (20°C approx.). Solubility in water: Fully soluble (20°C approx.).

Solubility in other solvents: Insoluble in organic solvents and lipids. Soluble in

surfactants, hydroalcoholic, hydroglyceric and hydroglycolic mixtures.

Heavy metals: No more than 5 mg/Kg. Sulfated ash: No more than 0.01%. Chlorides: No more than 0.007%. Sulfates: No more than 0.002%.

Other information: None.

Total aerobics: Max. 300 microorganisms / gram. Fungi and yeasts: Max. 200 microorganisms / gram.

Pathogens: Total absence in 1 gram.

DERMATOLOGY INNOCUOUSNESS

info@biogrundl.es

TECHNICAL DATA SHEET

It was performed by the patch test (Patch test) on 10 volunteers. To this purpose, 15% LINDEN EXTRACT by BIOGRÜNDL, S.L., was added to an O/W emulsion; 0,5g. of this emulsion was poured over the patch and attached on the forearm. It is thus maintained for 24, 48 and 72 hours, after which, the given reaction was evaluated, according to the following table of values:

- 1. No erythema.
- 2. Slight erythema.
- 3. Well defined erythema.
- 4. Severe erythema.

The result was absence of erythema on the 10 volunteers.

On the recommended concentrations, the product is skin innocuous.

STORAGE CONDITIONS

It is very important to keep the product in well closed containers, away from direct sunlight at temperatures not over 15°C. In case of not having refrigerators, store the product in the coolest and free from artificial light area.

INCI DENOMINATION

AQUA, GLYCERIN, TILIA CORDATA EXTRACT, POTASSIUM SORBATE.

EFFECTIVENESS TESTS

Assay: Double blind spot trial and subsequent cosmetic evaluation.

Assayed parameters: Emollient action. Calming action. Anti-inflammatory action.

N (volunteers): 30 (19 ♀ and 11 ♂). Age (volunteers): 19 - 59 years old.

Left side part to be treated:

Placebo, 30 volunteers, suitable cosmetic base.

Right side part to be treated:

LINDEN EXTRACT 2%, 15 volunteers, suitable cosmetic base.

Right side part to be treated:

LINDEN EXTRACT 5%, 15 volunteers, suitable cosmetic base.



TECHNICAL DATA SHEET


Assayed parameters:

	Assay	N (volunteers)	Age (volunteers)	Placebo	LINDEN EXTRACT 2%	LINDEN EXTRACT 5%	Assay performed (Days)	Results measured (Days)
Emollient action	Double blind spot trial and subsequent cosmetic evaluation.	N = 30 19♀ and 11 ♂	19 – 59	Left side part	Right side part 15 volunteers.	Right side part. 15 volunteers.	D = 30	D' = 1
Calming action	Double blind spot trial and subsequent cosmetic evaluation.	N = 30 19♀ and 11 ਕਾਂ	19 – 59	Left side part	Right side part 15 volunteers.	Right side part. 15 volunteers.	D = 30	D' = 30
Anti- inflammatory action	Double blind spot trial and subsequent cosmetic evaluation.	N = 30 19♀ and 11 ਕਾਂ	19 – 59	Left side part	Right side part 15 volunteers.	Right side part. 15 volunteers.	D = 30	D' = 1



TECHNICAL DATA SHEET

SKIN CONDITION EVALUATION AFTER 30 DAYS



(*) Emollient action was determined by softening degree of calluses on volunteers treated body zone.

Total number of days in which the assay was performed (treatment was applied): D = 30 days (every day during the effectiveness test).

Total number of days in which the results were measured: D' = 1 day (effectiveness test last day).

Average score:

Worst result: Zero limit value is assigned. **Best result:** Five limit value is assigned.

Score is assigned to obtained results between the two limit values.

Average computation for each obtained results group is done:

Placebo, 2% LINDEN EXTRACT by BIOGRÜNDL, S.L., 5% LINDEN EXTRACT by BIOGRÜNDL, S.L..

Results meaning:

AS rank between 0 - 1: No significant changes.

AS rank between 1 - 2: Slight improvement regarding placebo.

AS rank between 2 – 3: Appreciable improvement regarding placebo.


AS rank between 3 – 4: Great improvement regarding placebo.

AS rank between 4 – 5: Excellent cosmetic result.



TECHNICAL DATA SHEET

LINDEN EXTRACT 2%



Results:

Worst result LINDEN EXTRACT 2%: 18,5%. Average result LINDEN EXTRACT 2%: 29,5%. Best result LINDEN EXTRACT 2%: 41%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.


Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.



TECHNICAL DATA SHEET

LINDEN EXTRACT 5%



Results:

Worst result LINDEN EXTRACT 5%: 23%. Average result LINDEN EXTRACT 5%: 37%. Best result LINDEN EXTRACT 5%: 51%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.


Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.



TECHNICAL DATA SHEET

SKIN CONDITION EVALUATION AFTER 30 DAYS



(*) Calming action was measured by itch, redness and warm feeling decrease, by subjective assessment from volunteers, in the treated skin zone.

Total number of days in which the assay was performed (treatment was applied): D = 30 days (every day during the effectiveness test).

Total number of days in which the results were measured: D' = 30 days (every day during the effectiveness test).

Average score:

Worst result: Zero limit value is assigned. **Best result:** Five limit value is assigned.

Score is assigned to obtained results between the two limit values.

Average computation for each obtained results group is done:

Placebo, 2% LINDEN EXTRACT by BIOGRÜNDL, S.L., 5% LINDEN EXTRACT by BIOGRÜNDL, S.L..

Results meaning:

AS rank between 0 - 1: No significant changes.

AS rank between 1 - 2: Slight improvement regarding placebo.

AS rank between 2 – 3: Appreciable improvement regarding placebo.


AS rank between 3 - 4: Great improvement regarding placebo.

AS rank between 4 – 5: Excellent cosmetic result.



TECHNICAL DATA SHEET

LINDEN EXTRACT 2%



Results:

Worst result LINDEN EXTRACT 2%: 19%. Average result LINDEN EXTRACT 2%: 31%. Best result LINDEN EXTRACT 2%: 44%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.


Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.



TECHNICAL DATA SHEET

LINDEN EXTRACT 5%



Results:

Worst result LINDEN EXTRACT 5%: 29%. Average result LINDEN EXTRACT 5%: 41,5%. Best result LINDEN EXTRACT 5%: 54%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.


Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.



TECHNICAL DATA SHEET

SKIN CONDITION EVALUATION AFTER 30 DAYS



(*) Anti-inflammatory action was determined by direct measurement and observation of inflammation reduction from volunteers.

Total number of days in which the assay was performed (treatment was applied): D = 30 days (every day during the effectiveness test).

Total number of days in which the results were measured: D' = 1 day (effectiveness test last day).

Average score:

Worst result: Zero limit value is assigned. **Best result:** Five limit value is assigned.

Score is assigned to obtained results between the two limit values.

Average computation for each obtained results group is done:

Placebo, 2% LINDEN EXTRACT by BIOGRÜNDL, S.L., 5% LINDEN EXTRACT by BIOGRÜNDL, S.L..

Results meaning:

AS rank between 0 - 1: No significant changes.

AS rank between 1 - 2: Slight improvement regarding placebo.

AS rank between 2 - 3: Appreciable improvement regarding placebo.


AS rank between 3 - 4: Great improvement regarding placebo.

AS rank between 4 – 5: Excellent cosmetic result.



TECHNICAL DATA SHEET

LINDEN EXTRACT 2%



Results:

Worst result LINDEN EXTRACT 2%: 19%. Average result LINDEN EXTRACT 2%: 28,5%. Best result LINDEN EXTRACT 2%: 47,5%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.


Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.



TECHNICAL DATA SHEET

LINDEN EXTRACT 5%



Results:

Worst result LINDEN EXTRACT 5%: 27%. Average result LINDEN EXTRACT 5%: 43%. Best result LINDEN EXTRACT 5%: 59%.

Results meaning:

Al rank between < 20%: No significant changes.

Al rank between 21 – 40%: Slight improvement regarding placebo.

Al rank between 41 – 60%: Appreciable improvement regarding placebo.

Al rank between 61 – 80%: Great improvement regarding placebo.